

December 16, 2006

Dr. Raija Laukkanen
INWA Scientific Advisor

Review: Scientific evidence on Nordic Walking

This paper reviews 52 scientific articles published on Nordic Walking in years 1992-2006. Papers or results with at least English abstract are included. Studies are classified according to the target group into four categories:

1. Nordic Walking studies related to health and health-related fitness (done on sedentary, on elderly individuals or in patients)
2. Nordic Walking studies related to fitness (done on physically active individuals)
3. Nordic Walking studies related to sports type of performance (done on athletes)
4. Other (done on various groups, reviews, fitness assessment, safety and popularity of NW)

References of all studies are listed in alphabetical order.

1. Studies related to health

First research results on responses to pole walking training were published in 1992 by Stoughton, Larkin and Karavan from the University of Oregon. They studied psychological profiles (mood states) as well as muscular and aerobic fitness responses before and after 12 weeks of exerstriding or walking training in sedentary women. Exerstriding is a modified form of walking that incorporates the use of specially designed walking stick (Exertriders®) in a standard walking workout. This study group consisted of 86 20-50 year old women whose fitness was at moderate level. Maximal aerobic power (VO_{2max}) varied between 34-37 ml/kg/min. A study group was divided into three sub-groups. Control group did remain all their exercise habits. Walking and Exertrider groups did walk 30-45 minutes four times a week at the intensity corresponding to the 70-85 % maximum heart rate for twelve weeks. In Exertrider group both the walking speed and the distance walked were slightly less than in the walking group.

In both groups the maximal aerobic power and maximal treadmill time increased significantly. These increases were 8 and 19 % on an average. A slight increase in maximal ventilation occurred in the Exertrider group, but not in the walking group. Muscular endurance did improve in the Exertrider group by 37 % and in the walking group 14 %. Muscular strength assessed using triceps pushdown and a modified lateral pull-down did not improve in either group. Exertrider walkers showed significant improvements in depression, anger, vigor, fatigue, total mood disturbances

and total body-cathexis scores. The walking group showed significant improvements only in vigor and total body-cathexis. It was speculated that the Exertrider group may have felt more unique and special because of their opportunity to do a new and more enjoyable method of walking.

Exerstriders were also compared to the weighted vests, angle weights, hand and wrist weights, weighted gloves and Powerbelts™ by Porcari (1999) with similar results as above.

The effort involved during Nordic Walking has also been investigated in coronary heart patients (Walter et al. 1996). In this study fourteen men aged 61 years walked two eight minute repetitions: the first consisted of normal walking while during the second the subjects used half kilogram poles. The subjects walked at the maximum pace allowed in the light of their symptoms. All subjects had had either heart bypass or angioplasty operations or had suffered cardiac infarctions. During pole walking average energy consumption increased 21%, heart rate by 14 beats/minute and highest systolic/diastolic blood pressure figures by 16 and 4 mmHg respectively when compared with figures during normal walking. Oxygen pulse figures (i.e. oxygen consumption multiplied by heart rate) are indicative of changes in oxygen consumption and are not connected with undesirable rises in blood pressure. The research group concluded that pole walking is a safe form of rehabilitation for heart patients.

A Finnish study (Anttila et al. 1999) did compare Exel polewalking with the regular walking training for 12-weeks on 55 female office workers. The EMG-measurement showed that electrical activities of the muscles in the upper body neck-shoulder-upper back) were significantly higher when walking with poles. Polewalking training diminished neck and shoulder symptoms and subjective feeling of pain. Mobility of the upper body increased as well. The similar results were obtained also in the study by Karvonen et al. (2000). They did study neck-shoulder area pain on 31 44-50 year old person who had no previous experience on Nordic Walking. Exercise group did train twice a week for ten weeks and for 60 min per session. Nordic walking did decrease neck and shoulder pain in general and at work. In addition, the disturbance of neck and shoulder stiffness and pain in the movement of head were significantly decreased. A third study (Koskinen et al. 2003) did examine the effectiveness of NW on ageing employees on their postural control as well as to the muscular strenght of lower extremities and the middle trunk. Subjects (n=24) were 45 to 61 year old , majority of women. They did NW three times weekly and partly instructed. The group did improve in health-related fitness parameters evaluated by the Fitness Test battery developed by UKK Institute.

In a study by Baatile et al. (2000) 16 72-year old veteran males with Parkinson´s disease did Nordic Walk 8-weeks in an interval training program three times weekly for 60 min per session with the perceived rating of intensity (Borg 13). Researchers concluded that regular NW exercise program increased perceived functional independence and quality of life in individuals with Parkinson Disease.

The aim in Parkatti et al. study (2002) was to examine the benefits of NW on functions important to everyday life among older sedentary individuals in Virginia, USA. Altogether 18 73-year old (62-87 yrs) person participated. They did exercise

was 60 min twice a week for 12 weeks (10 min warm-up, stretching in the middle and cool-down at the end) by Nordic Walking. Functional capacity battery of tests included: chair stand, arm curl, chair sit and reach, scratch test, 2 min step in place and "up and go" test was used before/after intervention. Also a health questionnaire was used. The results of all functional tests were statistically significantly better after training. The study showed that NW is suitable for elderly and effective to affect on functional capacity.

In 2003 Collins et al. did publish a study where they studied 52 (65-70 yrs) patients with peripheral vascular diseases (PVD). The program consisted of three weekly sessions of pole walking for 30-45 min. Also Vitamin E (dose 400 IU daily) was studied. Pre- and post peak oxygen uptake, Quality of Life -interview and biweekly ankle blood pressure measurements were conducted. Pole group improved exercise tolerance significantly, they had also less claudication pain after exercise. Additionally, distance and walking speed improvement in pole groups. Researchers concluded that pole walking effectively improved both the exercise tolerance and perceived quality of life of patients with PVD. E-vitamin caused little additional benefit.

The same group did also publish another paper (Collins et al. 2004) in which the 24-week training data of 49 patients with peripheral arterial disease were studied. Persons with PAD were randomized into a 24-week PS training program or a non-exercise attention control (C) group. PS exercise entails walking with modified ski poles using a movement pattern similar to cross-country skiing. Those assigned to the PS group trained three times weekly. Control group subjects came to the laboratory biweekly for ankle blood pressure measurements, A symptom-limited ramp treadmill test, ratings of perceived leg pain, and the QOL data were obtained at baseline and upon completion of the training program. After 24-weeks of PS training, subjects increased their exercise endurance from 10.3 +/- 4.1 min to 15.1 +/- 4.5 min. This was significantly greater (7.46, $p < 0.001$) than C group subjects whose exercise endurance declined (11.2 +/- 4.7 to 10.3 +/- 4.7 min). Systolic blood pressure ($p < 0.001$), heart rate ($p = 0.04$), rate pressure product $p = 0.05$), oxygen uptake ($p = 0.016$) and perceived leg pain over exercise time on the baseline and six-month symptom-limited treadmill test were greater in the PS group when compared to the C group. The change in the physical component summary score of the SF-36 was also greater in the PS group when compared to the C group ($p = 0.03$). Conclusion was that in the training significantly improved the clinical indicators of increased cardiovascular fitness, improved the perception of the physical component of health-related QOL, and decreased symptoms of claudication pain during exertion in these patients.

Purpose of a research by Kukkonen-Harjula et al (2004) was to study training responses of brisk walking with or without poles on cardiorespiratory fitness (both in submaximal and maximal exercise) and on some other indicators of health-related fitness in healthy middle-aged women. Training prescription was aimed at moderate intensity (50-85 % of HR reserve, HRR). Two-hundred-and-twelve women volunteered. Their age was 50-60 years, no major health problems, BMI 20-30 kg x m⁻² and leisure exercise training no more often than twice weekly. After screening examinations 121 women were accepted and randomized into a NW or a walking (W) group. Walking and NW techniques instructed in small groups. Training intensity was

self-guided and resulted to be same in both groups (Mukka 2004). Training was 4x weekly for 40 mins, intensity 53% HRR, Borg 13.7. The increase in peak VO_2 (about 8 % in both groups) during 13 weeks' training was modest. Study also showed that the mode of maximal exercise testing (with or without poles) had no influence on peak VO_2 and its change during training in women with initially little familiarity with NW technique.

In a study by Aigner et al. (2004) 20 untrained healthy individuals (average age 47 years) were studied while walking with or without poles on separate days. The speeds of walks were 7.9 km/h on an average and mean heart rates 165 and 158 bpm with and without poles. The corresponding blood lactate levels were 5.7 and 5.0 mmol/l. In all speeds between 3 to 7 km/h heart rates and lactates were significantly higher in Nordic Walking compared to regular walking.

Heikkilä et al (unpublished 2004) did study NW in 13 33-54-year old overweight (BMI 32 kg/m²) individual. Subjects did exercise NW 4 months regularly HR controlled with progressive load. The results did show that subjects did lose weight (- 5kg on an average), and body fat (waist -6.6 cm), improved in aerobic fitness (+29%), and in blood lipids. The key factors, according researchers, to these very positive results were that intensity of training was moderate-high and that it was progressive in nature.

Wilk et al. (2005) did study NW in 16 acute coronary disease patients in Poland. Based on this exercise intervention they concluded that NW is a purposeful activity for cardiac rehabilitation.

Nordic walking has also been studied in cancer patients. In the recent study by Sprod et al. (2005) purpose was to determine the effects of walking pole use on shoulder function in female breast cancer survivors. Twelve 50-59 year old participants were randomly assigned into experimental or control groups for 8 weeks. Exercise program consisted of two 20 min aerobic training sessions at the intensity of 40-50% HRR. Additionally both groups did perform 30 min resistance training after aerobic session. No within group improvements were found in the control group. The experimental group did improve muscular endurance of the upper body. This was concluded to be beneficial in helping them to perform activities of daily living and regain an independent lifestyle.

In a study by Schöttler et al. (2005) Nordic Walking was studied for prevention in 150 patients with various orthopedic problems. They did exercise 30-90 min three times weekly with an educated instructor and many also at home by themselves. The results and experiences were asked by questionnaire. 93% felt that their endurance capacity did increase and in 55% body weight was reduced. Additionally, resistance to stress improved in 63% of patients and 21% felt that NW training did improve their sleep. The conclusions did agree with those from Heinz (2003).

Nordic Walking was related in type 2-diabetes patients by Nischwitz et al. (2006). Altogether 6 women and 13 males with the mean age of 67 years did participate NW 1 to 2 times weekly for 90 minutes for 1 year. Blood glucose values were taken at at stat and at 45 and 90 min during and 2 hours after walking sessions. Life quality questionnaires were filled at 3 months intervals. The overall results were very

encouraging indicating improvements in all diabetes-related metabolic parameters and did reduce the use of diabetes medication.

2. Studies related to fitness

The physiological responses to walking with and without Power Poles™ were studied by Hendrickson (1993) and by Porcari et al. (1997). Power Poles are specially constructed, rubber-tipped ski poles designed for use during walking. Hendrickson's study group consisted of sixteen fit women (VO_{2max} 50 ml/kg/min) and men (59). They did walk with and without poles on a treadmill with the speeds of 6-7,5 km/h. There were no differences in the responses between males and females. It was found that the use of poles significantly increased oxygen uptake, heart rate and energy expenditure by approximately 20 percent compared to the walking without poles in fit subjects. In Porcari's study on 32 healthy men and women walking with poles resulted in an average of 23% higher oxygen uptake, 22% higher caloric expenditure and 16% higher heart rate responses compared to walking without poles on a treadmill. RPE values averaged 1,5 units higher with the use of poles and the pattern of responses was similar for men and women.

A dual-motion treadmill Cross Walk has been studied by Knox (1993), Foley (1994) and by Butts et al. (1995). The Cross Walk Dual Motion Cross Trainer as a motorized treadmill designed to increase the energy cost of walking by incorporating arm activity during walking, thus increasing the muscle mass used during exercise. The activity is not the same as field walking with poles, but can be used as reference to NW. Knox did study thirty-seven 17-35 year old women and they all performed six 5-min steady-state exercises with and without arm activity. Walking with arm activity increased significantly heart rate, ventilation, oxygen uptake and energy expenditure compared to the walking without arm activity. E.g. heart rate increased 17-31 beats per minute. Rating of perceived exertion as well as energy expenditure increased with an average of 14 percent. In Butt's study both the 24-year old women and men were studied with a similar design. In this study arm work increased energy expenditure by 55 % on an average compared to the regular walking, but did increase rating of perceived exertion only little. This was consistent with the results from the Foley, who did study Cross Walk in 24-year-old men.

Rogers et al. (1995) did compare energy expenditure during submaximal walking with Exerstriders® in ten 24 year old fit women. Mean maximal aerobic power (21 vs. 18 ml/kg/min) and heart rate (133 vs. 122 bpm) were significantly greater during the walking with poles compared to walking without. Also the total caloric expenditure in a 30 minute session was significantly greater during pole walking (174 vs. 141 kcal). In contrast, the rating of perceived exertion did not differ significantly between the two conditions.

Laukkanen (1998, unpublished) did compare heart rate during normal and fast walking speeds with and without Exel Walker poles. Ten middle-aged men and women were studied on an indoor hall track. The heart rate increase, measured with

telemetric Polar HR monitor, was between 5-12 bpm and 5-17 bpm in women and men.

The purpose study by Knight and Caldwell (2000) was to compare pole and no-pole conditions during uphill backpacking, which was simulated on an inclined treadmill with a moderately heavy (22.4 kg, 30% body mass) backpack. Physiological measurements of oxygen consumption, heart rate, and RPE were taken during 1 h of backpacking in each condition, along with joint kinematic and electromyographic comparisons from data collected during a third test session. The results showed that although imposing no metabolic consequence, pole use elicited a longer stride length (1.27 vs 1.19 m), kinematics that were more similar to those of unloaded walking, and reduced activity in several lower extremity muscles. Although pole use evoked a greater heart rate (113.5 vs 107 bpm), subjects were backpacking more comfortably as indicated by their ratings of perceived exertion (10.8 vs 11.6). The increased cardiovascular demand was likely to support the greater muscular activity in the upper extremity, as was observed in triceps brachii. By redistributing some of the backpack effort, pole use alleviated some stress from the lower extremities and allowed a partial reversal of typical load-bearing strategies.

In a study by Jacobson et al (2000) load carriage energy expenditure with and without using hiking poles was studied. Twenty male volunteers aged 20-48yr completed two randomly treadmill trials with poles (E) and without poles (C). Poles and load (15 kg backpack) were fitted for each subject according to the manufacturers' suggestions. Heart rates (HR), minute ventilation (V(E)), oxygen consumption (O₂), caloric expenditure (Kcal), and rating of perceived exertion (RPE) were recorded at the end of each minute. Two trials separated by one week consisted of a constant treadmill speed of 1.5 mph and 1 min at 10% grade, 2 min at 15% grade, 2 min at 20% grade, and 10 min. at 25% grade. Analysis of paired time points yielded no significant differences in HR, VO₂, V(E), and Kcal, however, RPE means were significantly lower for 5 of the last 7 trial minutes with the use of poles. These results suggest that during load carriage on moderate grade, the weight and use of hiking poles does not increase energy expenditure but may provide reduced perceptions of physical exertion.

Gullstrand & Svedenhag (2001) from Sweden did study acute physiological effects on walking on a treadmill with or without poles. This study on 13 55-year old subjects did show that VO_{2max}, VE, blood lactate and HR did increase, but RPE (rating of perceived exertion) remained unchanged in NW compared to regular walking

In the study published by the Cooper Institute group from Texas, USA the metabolic cost of NW was compared to normal walking in 22 31-year-old men and women (Morss et al. 2001, Church et al. 2002). Participants of this study did walk on an outdoor 200-m track with Cosmed K4b for oxygen consumption and Polar Vantage heart rate monitor for HR measurements. Study indicated significant increases of oxygen consumption (20% on average), caloric expenditure and HR in NW compared to normal walking. The range of increase was large, i.e. in oxygen consumption 5-63% indicating differences in poling intensity and technique. Perceived exertion did not differ between the walks. Same group did also compare separately metabolic cost

of high intensity poling (Jordan et al. 2001). In high intensity poling NW increased HR 35 bpm on an average compared to regular walking.

In a study by Willson et al. (2001) the purpose was to determine whether walking with poles reduces loading to the lower extremity during level over ground walking. Three-dimensional gait analysis was conducted on 13 healthy adults who completed 10 walking trials using three different poling conditions (selected poles, poles back, and poles front) and without the use of poles (no poles). Results did show that there were differences in kinetic variables between walking with and without poles. The use of walking poles enabled subjects to walk at a faster speed with reduced vertical ground reaction forces, vertical knee joint reaction forces, and reduction in the knee extensor angular impulse and support moment, depending on the poling condition used.

Bohne study (2002) did study poles use effectiveness during downhill hiking in 20 men and women. Gender differences were found indicating that men did decrease ground reaction forces with poles at steeper hill (25°) than women (20°) did.

A study done in Germany by Ripatti (2002) 24 individuals (48±8 yrs) did NW for 6 weeks 2 times weekly for 60 min (65-85 %HRmax). This improved their endurance capacity even walking at lower speed.

Parker et al. (2002) did compare metabolic responses to graded exercise walking with and without poles in 14 physically active 30-year old subjects. The comparisons of heart rate and metabolic parameters were done in laboratory on treadmill and using cross-country skiing poles. The tests did not show any difference between the exercise modes.

Effects of regular Nordic Walking training in elderly (mean age 76 years) men and women was studied by Luoma-aho (2002). Subjects did walk 3 times 50 minutes weekly for two months. The performance in six-minute test improved 14%, maximal walking speed 20%, and balance 4%. Health-rated health improved 22%, self-rated physical fitness 24% and performance in the physical activities of daily living 30%.

Mänttari et al (2004) did conduct a pilot study for Kukkonen-Harjula et al. intervention study (2004). In this pilot they compared the cardiorespiratory and musculoskeletal responses of NW and W in field conditions in middle-aged women, with three self-guided exercise intensities. After screening examinations 20 middle-aged women performed a maximal exercise test on a treadmill with poles. All the subjects were familiar with Nordic walking or cross-country skiing. These results showed that Nordic walking increased the mean HR compared to regular walking only from 2.6% to 4.9% and the mean VO_2 from 2.5% to 10.8%, during the three different self-guided walking intensities. This increase seems to be due to the increased muscle activity in the upper body muscle groups. Compared to previous studies the statistically significant mean differences between NW and W were modest.

In Kleindienst et al. study (2006) the biomechanical differences between the locomotion patterns of NW, walking and running were studied in eleven subjects experienced in NW. High speed camera systems picturing simultaneous posterior and

lateral movements were used. The data reveal differences between the three analysed locomotion patterns. For NW as well as walking the mechanical load of the lower extremity was lower compared to running. None of the kinematic parameters suggested a "physiological benefit" of NW compared to walking. Moreover NW shows higher vertical and horizontal forces during landing. Exclusively the lower vertical force peak during push off indicated a lower mechanical load for NW in comparison to walking. The authors did conclude that it is questionable if NW - based on its promised "biomechanical benefits" compared to walking - should be recommended for overweight people and for people with existing musculoskeletal problems of the lower limb.

In a study by Schiffer (2006) the physiological responses during incremental field tests (FT) in Nordic walking (NW), walking (W) and jogging (J) were studied. Fifteen healthy middle-aged women participated in three FT. Heart rate (HR) and oxygen uptake were monitored continuously by portable analysers. As a main result and conclusion the authors did state that as NW speed at submaximal lactate levels is lower than in W and J, W and J test measures of HR are not suitable for NW training recommendations.

3. Studies related to sports

In a Norwegian study by Haugan and Sollesnes (2003) 16 sports students (22 yrs) were measured in a laboratory walking at the speeds of 5.5, 6.0 and 6.5 km/h with or without poles on an elevated treadmill (17%). Half of the subjects were cross-country skiers. Oxygen uptake increased significantly at all speeds when using poles in walking in others, but not in c-c skiers.

4. Other

Nordic Walking has also been under study in The Netherlands. Lande et al. did publish in 2003 a systematic review of the physiological effects of pole walking.

Parkkari et al. (2004) did evaluate injury risk in various commuting and lifestyle activities in a cohort of 3657 15-74-year old Finns. The individual injury risk per exposure time was overall relatively low, ranging from 0.19 to 1.5 per 1000 hours of participation. Highest risk in all recreational and competitive sports was in squash (18.3), judo (16.3) and orienteering (13.6). In Nordic Walking (pole walking) the risk was 1.7. In the cohort 11 % participated actively this sports.

Knobloch and Vogt (2006) did study safety of NW training prospectively by questionnaires in 137 persons with the mean age of 53 years. Injury rate was 0.93/1000 hours of training. Most frequent injury was distorsion of the ulnar ligament of the thumb, following with the shoulder and gastrocnemius muscle injuries. Recovering back to normal training took place in 4 weeks. Overall conclusion of this large scale study was that NW is safe.

UKK Walk Test is a popular test for aerobic fitness assessments in large population surveys since it is easy, safe and cost effective. The Test has been developed and is aimed to be done with regular walking style. Widmann et al. (2005) did study the

possible differences in conducting the 2-km UKK test with regular walking style or with nordic walking. Altogether 17 women (mean age 55 years) did participate into Test with and without poles in two different days. The results did not show any differences in walking time and in physiological parameters (heart rate, respiratory quotient, oxygen uptake, blood lactate, energy expenditure). The result was same in those who had good technique compared to less skilled ones.

In a questionnaire study by Schmidt et al. (2004) 226 German adults (66% women) who practiced NW regularly were interviewed during winter 2003-2004. The average age was 52 yrs and BMI 25 kg/m². The main motivation for NW was health, 12% wanted to test something new, 6% did it as an option for c-c skiing in summer. 71% worked out for arm and trunk muscles, 23% in order to reduce joint load. 54% would prefer to have a similar net of trails like those for hiking.

Nordic Walking has also been studied from the consumer perspective (Shove and Pantzar 2004). Authors conclude that popularity of the NW has arisen through the active and ongoing interaction of images, artifacts and forms of competence; a process in which both consumers and producers are both involved.

Popularity of NW has been studied in Finland in two studies (National Exercise Study 2005-2006 by Suomen Kuntourheiluliitto, Nordic Walking survey 2006 in Finland by Suomen Latu). In National study phone calls interview based gallup-survey was done in 5500 Finns did indicate NW being the second most popular activity mode in Finnish adults. Walking is the most popular (1 840 000 exercisers) activity. Altogether 444 000 men and women of 19-65-year old adults do nordic walk regularly in Finland 2005-2006. The increase in nordic walkers in Finland since 2001-2002 was +144 000 adults (+48%), which places NW the second biggest activity mode in grow.

In a NW survey 2006 published by Suomen Latu association 1000 over 15 year old Finns were interviewed. Participation in NW in Finland has been studied since 1998. According 2006 results the number of people experiencing Nordic Walking in Finland seems to remain in 1,5 million. 720 00 Finns do NW at least once a week; of them 39% over 65-year old. In 2006 data 26% of adult women and 11% of adult men NW regularly. The numbers indicate that NW has become more popular among men.

The different results of these two studies are explained by different form of questionnaires used.

Summary

To summarize the acute physiological effects of Nordic walking, it increases the energy consumption of the body compared to regular walking with the same speed without poles both in women and men of all ages and in fit and less fit individuals. The increase is due to larger working muscle mass in the upper body. The increase varies individually according to walking speed and technique. If the speed is very fast, there is less time for efficient pushing off with poles and thus decreased upper body muscular involvement. Similarly to energy consumption the increase in heart rate is variable. Because perceived exertion in pole walking is often less than true

physiological strain, controlling heart rate may be beneficial for those tending to overreach. The resulting increases in energy consumption and heart rate in Nordic walking mean that the cardiovascular strain induced by Nordic walking is greater compared to walking without poles at the same speed. This is desirable for those people who have difficulty reaching their training heart rate by walking - instead of having to start running they can start using walking poles and continue walking. Even studies show some disagreement on biomechanical benefits of NW, it involves less harmful impacts to the lower extremities compared to running, and therefore may prevent from injuries.

To summarize, the training effects of Nordic walking on cardiorespiratory fitness and endurance have been shown to be similar to walking training in middle-aged and elderly women. In fit individuals and in men intervention studies are missing. In the studies, the improvement in Nordic walking was reached by lower speed and thus by shorter distance walked, because the cardiovascular strain was greater in Nordic walking than in ordinary walking without poles if the same speed was used. Walking with poles improves mainly aerobic fitness, muscular endurance, decrease neck-shoulder area disabilities and pain and can have positive effects on mood state. In order to improve muscle power, uphill walking is required. Pole walking affecting on body coordination and motor fitness has been published little. Nordic Walking is safe activity and individuals are motivated to Nordic Walk mainly due health reasons.

Even there is rather strong scientific evidence on both acute and long-term effects of Nordic Walking some research challenges still remain. Randomized controlled trials on dose-responses of health and fitness improvements in men and in women, in healthy, in fit and in individuals with minor health problems (body weight, insulin-resistance, blood pressure, osteoporosis) are still lacking. Additional biomechanical studies on NW effects on musculoskeletal health are needed. Also, studies in motivation and adherence in NW as well as overall global participation (walkers, their demographics and their social and other status) in this activity is missing.

Author:

Dr. Raija Laukkanen
 Ph.D, FACSM
 Docent, University of Oulu, Department of Medicine
 Director, Exercise Science
 Polar Electro Oy
 Professorintie 5
 90440 Kempele
 Finland
 Tel.+358 8 5202100, GSM +358 400 588624
 Fax.+358 8 5202331
 e-mail raija.laukkanen@polar.fi

For more information and original studies contact:

Ms. Tiia Aherto
 Association secretary
 International Nordic Walking Association

Valimotie 5
 FIN-01510 Vantaa
 Finland
 tiia@inwa.nordicwalking.com

INWA web home page:

www.inwa.nordicwalking.com

References

Aigner A, Ledl-Kurkowski E, Hörl S, Salzmann K. Effecte von Nordic Walking bzw. Normalen Gehen auf Herzfrequenz und arterielle Laktatkonzentration. Österreichisches Journal für Sportmedizin 34, H.3, 32-36, 2004.

Anttila, Holopainen, Jokinen. Polewalking and the effect of regular 12-week polewalking exercise on neck and shoulder symptoms, the mobility of the cervical and thoracic spine and aerobic capacity. Final project work for the Helsinki IV College for health care professionals, 1999.

Baatile J, Langbein W, Weaver F, Maloney C, Jost M. Effect of exercise on perceived quality of life of individuals with Parkinson's disease. J of Rehabilitation Research and Development 37(5), September/October 2000.

Bohne MJ Effects of hiking downhill with poles on selected kinematic and kinetic parameters. Thesis at Utah State University 2002.

Butts, Knox, Foley. Energy cost of walking on a dual-action treadmill in men and women. Med Sci Sports Exerc 27(1), 121-125, 1995.

Church, Earnest, Morss. Field testing of physiological responses associated with Nordic Walking. Res Quart Exerc Sports 73;296-300, 2002.

Collins E, Langbein W, Orebaugh C, Bammert C, Hanson K, Reda D, Edwards L, Littooy F. PoleStriding exercise and vitamin E for management of peripheral vascular disease. Med Sci Sports Exerc 3;384-393, 2003.

Collins EG, Langbein WE, Orebaugh C, Bammert C, Hanson K, Reda D, Edwards L, Littooy F N Cardiovascular training effect associated with polestriding exercise in patients with peripheral arterial disease Circulation 110 (17, Suppl. S): p 665, 2004.

Foley. The effects of Cross Walk (R)'s resistive arm poles on the metabolic costs of treadmill walking. Thesis. University of Wisconsin-La Grosse, 1994.

Gullstrand L, Svedenhag J. Training effects after 7 weeks of pole- and normal walking. 8th Annual Congress of the European College of Sport Science. Salzburg, Austria 09.-12.07.2003. In: Abstract book. Ed. Mueller E, Schwameder H, Zallinger G, Fastenbauer V. Institute of Sport Science, University of Salzburg, Austria 2003, p. 33-34 (abstract).

Haugan A, Sollesnes B. Does submaximal oxygen uptake increase when using Nordic Walking poles? Academic degree study. Sogn og Fjordane University College Faculty of Teacher Education, Sogndal, Norway, May 2003.

Heikkilä M, Kettunen O, Vasankari T. Improved fitness and reduced weight with Nordic Walking. Report. Vierumäki, Finland, December 2004, unpublished.

Heinz M. Nordic Walking. Auch als Rehabilitationsmethode geeignet. Nordic Walking. Also suitable in rehabilitation. Sportverletzung Sportschaden: Organ der Gesellschaft für Orthopädisch-Traumatologische Sportmedizin, 2003, vol 17(2); p.55.

Lande S, Hermansen B, Stamland O, Nordaune P. The physiological – and additional effects of pole walking. A systematic review. Research report at Fontys University of Professional Education Eindhoven, Department of Physiotherapy. Eindhoven, The Netherlands, 2003.

Hendrickson. The physiological responses to walking with and without Power Poles™ on treadmill exercise. Thesis. University of Wisconsin-La Grosse, 1993.

Jacobson B H, Wright T, Dugan B J. Load carriage energy expenditure with and without hiking poles during inclined walking. International journal of sports medicine, 21 (5),356-9, July 2000.

Jordan, Olson, Earnest, Morss, Church. Metabolic cost of high intensity poling while Nordic Walking versus normal walking. Med Sci Sports Exerc 33(5), S86, 2001.

Karawan. The effects of twelve weeks of walking or Exertriding on upper body muscular strength and endurance. Thesis. University of Wisconsin-La Grosse, 1992.

Karvonen, Mörsky, Tolppala, Varis. The effects of stick walking on neck and shoulder pain in office workers. Final project work at Mikkeli Polytechnic School. Degree programme of Physiotherapy. Mikkeli 2001.

Kleindienst FI et al. Comparison of kinematic and kinetic parameters between the locomotion patterns in nordic walking, walking and running. Sportverletzung Sportschaden- Organ der Gesellschaft für Orthopädisch-Traumtologische Sportmedizin, 20(1),25-30, March 2006.

Knight CA, Caldwell GE. Muscular and metabolic costs of uphill backpacking: are hiking poles beneficial? Med Sci Sports Exerc, 32 (12) 2093-101. Dec 2000.

Knobloch K, Vogt PM. Nordic Walking Injuries - Nordic Walking Thumb as Novel Injury Entity. Sportverletz Sportschaden 20: 137-142, 2006.

Knox. Energy cost of walking with and without arm activity on the Cross Walk dual motion cross trainer. Thesis. University of Wisconsin-La Grosse, 1993.

Koskinen J, Kärki M, Virtanen M. Power and Balance from Nordic Walking – effects of regular NW to muscular strenght and postural control of ageing employees who are

unaccustomed to regular physical exercise. Bachelor of Physiotherapy degree. Helsinki Polytechnic Health Care and Social Services, Helsinki, 2003.

Kukkonen-Harjula K, Mänttari A, Hiilloskorpi H, Pasanen M, Laukkanen R, Suni J, Fogelholm M, Parkkari J. Training responses of self-guided brisk walking with or without poles- a randomized controlled trial in middle-aged women. 9th Annual Congress of the European College of Sport Science. Clermont- Ferrand, France, 03.-06.07.2004. Book of abstracts, p.157.

Langbein WE, Collins EG, Orebaugh C et al. Increasing exercise tolerance of persons limited by claudication pain using polestriding. *J Vasc Surg* 2002;35:887-893.

Laukkanen R. HR response to Nordic Walking compared to regular walking. 1998 (unpublished).

Larkin. Aerobic responses to 12 weeks of exercitriding or walking training in sedentary adult women. Thesis University of Wisconsin-La Grosse, 1992.

Luoma-Aho Marika. Effects of supervised Nordic Walking on physical performance in elderly men and women living in sheltered houses. Department of Health Sciences. University of Jyväskylä.2002.

Morss, Church, Earnest, Jordan. Field test comparing the metabolic cost of normal walking versus walking with Nordic Walking. *Med Sci Sports Exerc* 33(5), S23, 2001.

Mukka M. Estimating of self-guided Nordic Walking and walking training intensity in 50-60 years old women. University of Jyväskylä, faculty of sports and health sciences, Department of health sciences. 2004.

Mänttari A, Hannola H, Laukkanen R, Hiilloskorpi H, Alikoski J, Valve R, Pekkarinen H, Parkkari J. Cardiorespiratory and musculoskeletal responses of walking with and without poles in field conditions in middle-aged women. 9th Annual Congress of the European College of Sport Science. Clermont- Ferrand, France, 03.-06.07.2004. Book of abstracts, p. 157

National Exercise Study 2005-2006. Suomen Kuntourheiluliitto, Kunto ry., SLU:n julkaisusarja 04/06, Helsinki 2006 (www.kunto.fi).

Nischwitz M, Meier GR, Dieterle C et al. Nordic Walking bei Patienten mit Diabetes mellitus Typ 2. *Diabelologie und Stoffwechsel* 2006;1.

Nordic Walking survey 2006 in Finland. Suomen Latu (www.suomenlatu.fi).

Parkatti T, Wacker P, Andrews N. Functional capacity from Nordic Walking among elderly people. Seminar poster at University of Jyväskylä, Finland, 2002.

Parker DL, Quintana R, DeWitt R. Metabolic responses to graded exercise walking with and without poles. *Med Sci Sports and Exerc*, 34 (5 Suppl), S295, May 2002

Parkkari J, Kannus P, Natri A, Lapinleimu I, Palvanen M, Heiskanen M, Vuori I, Järvinen M. Active living and Injury Risk. *International Journal of Sports Medicine* 25:209-216, 2004.

Porcari, Hendrickson, Walter, Terry, Walsko. The physiological responses to walking with and without Power Poles™ on treadmill exercise. *Res Quart Exerc Sports* 68(2),161-166,1997.

Porcari. Pump up your walk. *ACSM's Health and Fitness Journal* Jan/Feb,25-29, 1999.

Ripatti T. Effect of Nordic Walkin training program on cardiovascular fitness. Academic degree study. *Sportartspezifische Leistungsfähigkeit* Deutsche Sporthochschule Köln, Germany, 2002.

Rodgers, Vanheest, Schachter. Energy expenditure during submaximal walking with Exerctriders®. *Med Sci Sports Exerc* 27(4), 607-611,1995.

Schmidt A, Sonnen J, Krämer A, Roth R. Nordic Walking- an analysis of target groups and percpectives. 9th Annual Congress of the European College of Sport Science. Clermont-Ferrand, France, 03.-06.07.2004. Book of abstracts.

Schiffer T et al. Physiological responses to nordic walking, walking and jogging. *Eur J Appl Physiol*. Jun 24, 2006.

Schöttöer M, Kuhn W, Frey A. Nordic Walking – eine moderne alltagstaugliche form der prävention? *Phys Rehab Kur Med* 2005;15.

Shove E, Pantzar M. Consumers, producers and practices: understanding the invention and reinvention of Nordic Walking. *Journal of Consumer Culture* (in press 2004).

Sprod LK, Drum SN, Bentz AT, Carter SD, Schneider CM. The effects of walking poles on shoulder function in breast cancer survivors. *Integrative Cancer Therapies* 4(4);287-293, 2005.

Stoughton. Psychological profiles before and after 12 weeks of walking or Exertrider training in adult women. Thesis. University of Wisconsin-La Grosse, 1992.

Walter, Porcari, Brice, Terry. Acute responses to using walking poles in patients with coronary artery disease. *J Cardiopulm Rehabil* Jul-Aug;16(4), 245-50, 1996.

Widmann N, Haertel S, Bös K. Stress parameters at the UKK 2-km Walk Test with and without walking poles. Staatsexamenarbeit at University of Karlsruhe, 2005.

Wilk M, Kocur P, Przywarska I, Rozanska A, Owczarski T, Dylewicz P, 2005 (unpublished)

Willson J, Torry M, Decker M, Kernozek T, Steadman J. Effects of walking poles on lower extremity gait mechanics. *Journal: Medicine and Science in Sports and Exercise* 33 (1): p 142-147, January, 2001.

www.benidormnordicwalking.es